

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Stellar disc – dynamical evolution in a perturbed potential

Ladislav Šubr

Astronomical Institute, Charles University in Prague

Abstract. Models of the origin of young stars in the Galactic Centre are facing various problems. The most promising scenario of the star formation in a thin self-gravitating disc naturally forms stars on coherently rotating orbits, but it fails to explain origin of several tens of stars that evidently do not belong to any of the disc-like structures in the GC. One possible solution lies in rather complicated initial conditions, assuming at least two infalling and interacting gas clouds. We present alternative solution showing that a single thin stellar disc may have given birth to all young stars in the GC. The outliers are explained as stars that have been stripped from the parent structure due to the gravitational interaction with the gaseous circum-nuclear disc.

We investigate a model of the Galactic Centre which contains following constituents: **Supermassive black hole (SMBH)** of mass $M_{\bullet} = 3.5 \times 10^6 M_{\odot}$ dominates the gravitational field and it is approximated by a fixed Keplerian potential. **Circum-nuclear Disc (CND)** is modelled by several tens of particles orbiting the SMBH in a toroidal structure. Its total mass is not well determined by observations; we assume $M_{\text{CND}} \approx 0.2 M_{\bullet}$ orbiting at a characteristic radius of $R_{\text{CND}} \approx 1.5 \text{ pc}$. **Stellar cusp** of mass $\gtrsim 0.1 M_{\bullet}$ within 1 pc is assumed to be spherically symmetric. We incorporate it by means of a smooth fixed potential. **Young stellar disc** is a set of ~ 100 equal-mass gravitating particles. Initially, they are on circular orbits with nearly colinear angular momenta. Semi-major axes have distribution $\propto a^{-1}$ within the interval $(0.04 \text{ pc}, 0.4 \text{ pc})$.

Provided the CND would be the only perturbation to the SMBH's potential, individual stellar orbits would undergo secular evolution. In particular, orbital eccentricity and inclination would oscillate on the time-scale comparable to the age of the young stars. These oscillations are considerably damped if additional *spherically* symmetric perturbation of the late type stellar cusp is present. This component, however, does not suppress differential precession of the stellar orbits around the symmetry axis of the CND. The precession rate can be approximated as Šubr et al. (2009)

$$\frac{d\Omega}{dt} \approx -\frac{3}{4} \cos i \frac{M_{\text{CND}}}{M_{\bullet}} \frac{a\sqrt{GM_{\bullet}a}}{R_{\text{CND}}^3} \frac{1 + \frac{3}{2}e^2}{\sqrt{1 - e^2}} \approx \text{const}.$$

There is a strong dependence of $d\Omega/dt$ on the semi-major axis and the orbit inclination i with respect to the plane of the CND. Stars at the outer part of the disc and/or stars that are less perpendicular to the CND undergo faster precession, i.e. the disc becomes warped. The shape of the stellar disc in terms of the normal vectors of the orbits after several Myrs of such a differential precession is shown in the right panel of Fig. 1. We have verified Šubr et al. (2009) that such a model of a warped stellar disc is compatible with the observational




Figure 1. Dissolution of the coherently rotating structure due to the gravity of stellar cusp and the CND in terms of the direction of the angular momenta. Self-gravity of the stellar disc is considered only in the right panel, while it is omitted in the left one. Normal vectors of all orbits lie within the shaded area at $T = 0$; crosses represent their position at $T = 7\text{Myr}$. Coordinate system is aligned with the axis of the CND.

data available in the literature Paumard et al. (2006); Bartko et al. (2009). The picture changes when self-gravity of the stellar disc is taken into account (see Fig. 1). It appears that mutual torques of the stellar orbits tend to drive the coherently rotating ‘core’ of the disc towards the orientation perpendicular to the CND. At the same time, two-body relaxation scatters some stars to orbits that undergo fast precession and do not apparently belong to the parent structure after few Myrs. Another consequence of the gravitational interaction of the stellar orbits with both the CND and stellar disc itself is an enhanced growth of orbital eccentricities Haas et al. (2010).

Conclusions

We suggest that all young stars (probably except for S-stars) in the Galactic Centre could have been formed in a single thin self-gravitating disc. Perturbative influence of the CND then led to partial destruction of the coherently rotating structure. In addition to our previous results, we have found that the core of the disc has tendency to migrate towards orientation perpendicular to the CND, which is in a striking accord with the data provided by the observations. This result cannot be achieved if the gravity of either the CND, the spherical cusp or the stellar disc is ignored.

Acknowledgments. This work was supported by the Czech Science Foundation (ref. 205/07/0052) and the Centre for Theoretical Astrophysics in Prague.

References

- Bartko H., Martins F., Fritz T. K., et al., 2009, ApJ 697, 1741
- Haas J., Šubr L., Kroupa P., 2010, in preparation
- Paumard T., Genzel R., Martins F. et al., 2006, ApJ 643, 1011
- Šubr L., Schovancová J., Kroupa P., 2009, A&A 496, 695